

gctgctcagc tggaaagaa ggtggctgtg gctgactatg tggAACCTC tccccaggc	240
accaagtgg gccttgggg cacatgtgc aacgtgggt gcataccaa gaagctgatg	300
catcaggctg cactgtggg gggcatgatc agagatgctc accactatgg ctgggaggtg	360
gcccagcctg tccaaacacaa ctggaagaca atggcagaag ccgtgaaaa ccatgtgaaa	420
tccttgaact ggggtcattcg cgtccaaactg caggacaggaa aagtcaagta cttaacatc	480
aaagccagct ttgtggatga gcacacagtt cgccgtgtgg acaaaggcgg gaaggcgact	540
ctgcttcag ctgagcacat tgcattgtc acaggaggac ggccaaggta ccccacacaa	600
gtcaaaggag cccttggata tggaatcaca agtgacgaca tcttctggct gaaggagtc	660
cctggaaaaa cgttgggtgg tggagccagc tatgtggccc tagagtgtgc tggcttcctc	720
actggaaattt gactggatac cactgtcatg atgcgcgacca tccctctccg aggcttgac	780
cagcaaattgt catcttttgtt cacagagcac atggagtctc atggcacccca gttcctgaaa	840
ggctgtgtcc cctccacat caaaaaactc ccaactaacc agctgcaggt cacttggag	900
gatcatgctt ctggcaagga agacacaggc acctttgaca ctgtcctgtg ggccatagg	960
cgagttccag aaaccaggac tttgaatctg gagaaggctg gcatcgtac caaccctaag	1020
aatcagaaga ttattgtgg tggccaggag gctacctctg ttccccacat ctatgccatt	1080
ggagatgttgc tgagggggcg gcctgagctg acgcccacag ctatcaaggc aggaaagctt	1140
ctggctcagc ggcttttgtt gaaatcctca accttaatgg attacagcaa tgcccaca	1200
actgtcttta caccacttgg gtagtggctgt gtggggctgt ctgaggaggaa ggctgtggct	1260
ctccatggcc aggagcatgt agaggtttac catgcattt ataagccct agagttcagc	1320
gtggcggata gggatgcac acagtgcacata aaaaatgg tgcagatgg ggagccccca	1380
caacttggc tggccttgc cttccttggc ccaacgcgtg gagaagtcac ccaaggattt	1440
gctcttggaa tcaagtgtgg ggcttcatat gcacagggtga tgcagacagt agggatccat	1500
ccacactgt ctgaggagggt ggtcaagctg cacaatctca agcgctccgg cctggagct	1560
actgtgactg gttgtgtca	1578

## What is claimed is:

1. A method for the prevention of the reduction of a disulfide bond in an antibody expressed in a recombinant host cell, comprising, following fermentation, sparging the pre-harvest or harvested culture fluid of said recombinant host cell with air, wherein the amount of dissolved oxygen ( $dO_2$ ) in the pre-harvest or harvested culture fluid is at least 10%.

2. The method of claim 1 wherein said air sparging is continued until the pre-harvest or harvested culture fluid is at least 30% saturated with air.

3. The method of claim 1 wherein said air sparging is continued until the pre-harvest or harvested culture fluid is between about 100% saturated to about 30% saturated with air.

4. The method of claim 1 wherein the amount of dissolved oxygen ( $dO_2$ ) in the pre-harvest or harvested culture fluid is at least 30%.

5. The method of claim 1 wherein the antibody is a therapeutic antibody.

6. The method of claim 1 wherein the antibody is a biologically functional fragment of an antibody.

7. The method of claim 1 wherein the host cell is eukaryotic host cell.

8. The method of claim 7 wherein the eukaryotic host cell is a mammalian host cell.

9. The method of claim 1 wherein the host cell is prokaryotic host cell.

10. The method of claim 9 wherein the prokaryotic host cell is a bacterial cell.